

Complement Enhances the Pathogenesis in Autoinflammation Via Induction of A Pro-Inflammatory Microenvironment

Brunner J^{1*}, Posch W², Rabensteiner E¹ and Wilflingseder D²

¹Department of Pediatrics and Institute of Hygiene, Medical University Innsbruck, Austria

²Department of Medical Microbiology, Medical University Innsbruck, Austria

*Corresponding author:

Juergen Brunner,
Department of Pediatrics, Medical University
Innsbruck, Anichstrasse 35, 6020 Innsbruck,
Austria, Tel: 0043 512 504 80870,
E-mail: Juergen.brunner@tirol-kliniken.at

Received: 03 July 2021

Accepted: 22 July 2021

Published: 27 July 2021

Copyright:

©2021 Brunner J. This is an open access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and build upon your work non-commercially.

Citation:

Brunner J, Complement Enhances the Pathogenesis in Autoinflammation Via Induction of A Pro-Inflammatory Microenvironment. *Ann Clin Med Case Rep.* 2021; V7(3): 1-2

Keywords:

Autoimmune diseases; Autoinflammation; Inflammation; Interleukins; Pediatric rheumatic diseases

1. Introduction

The complement system is a complex part of the innate immune system and consists of more than thirty different proteins present in the plasma and on cell membranes. Complement can be activated via three different pathways and may play a role in triggering the NLRP3 inflammasome [1]. Inflammasomes are essential components in innate immunity. A group of inflammasome disorders has been associated with Autoinflammatory Diseases (AIDs). Familial Cold Autoinflammatory Syndrome (FCAS), Muckle-Wells Syndrome (MWS), and Chronic Infantile Neurological, Cutaneous and Articular syndrome/Neonatal Onset Multisystem Inflammatory Disease (CINCA/NOMID) were originally described as three distinct conditions. After the identification of their common genetic origin, i.e., mutations in the NLRP3 gene on chromosome 1q44, they are categorized as a continuum of one disease entity and labeled Cryopyrin-Associated Periodic Syndromes (CAPS). This preliminary study on a patient with MWS aimed to investigate the interplay between the complement system and this autoinflammatory disorder. PBMCs (Peripheral Blood Mononuclear Cells) were isolated from blood of a healthy donor and an MWS patient by density gradient centrifugation using a Ficoll Paque Premium (GE Healthcare). After washing, PBMCs were incubated with anti-human CD14 Magnetic Beads (BD) to obtain CD14⁺ monocytes. These were stimulated with cytokines (IL-4 and GM-CSF) for five days to generate immature monocyte-derived DCs (iDCs). iDCs were stimulated with either LPS or differentially opsonized HIV-1

preparations (non-opsonized HIV-1, HIV; complement-opsonized HIV-1, HIV-C). After two days in culture, iDCs and low and high LPS concentrations (10 ng/ml and 100 ng/ml) or HIV-exposed DCs were used for cytokine ELISAs and flow cytometric analyses. IL-1 β and IL-6 ELISAs were analyzed and phenotypical characterization of pathogen-exposed DCs was performed by analyzing characteristic surface markers (CD11c, DC-SIGN, CD86) by multi color flow cytometry [3]

2. Results

IL-1 β and IL-6 production of iDCs in the patients' cells was significantly higher upon stimulation with non- and complement-opsonized HIV-1, with complement-opsonized HIV-1 being the most potent stimulator (Figure 1a and 1b). In contrast, the bacterial stimulus LPS mediated a non-significant increase in IL-1 β at the high concentration in the individual suffering from MWS, while IL-6 was only slightly upregulated by the higher LPS concentration in both the healthy and MWS individual (Figure 1a and 1b). We recently described CD11c, the alpha chain of CR4, as a trigger of pro-inflammatory cytokines and type I interferon production [4]. From these findings we conclude that the complement system may play an essential role in the development of an overshooting pro-inflammatory milieu in patients with autoinflammatory disorders. The phenomenon shown in a patient with MWS should be reproduced in more MWS patients as well as in patients with other autoinflammatory disorders.




Figure 1a: IL-1β and IL-6 production




Figure 1b: IL-1β and IL-6 production of iDCs in the patients’ cells was significantly higher upon stimulation with non- and complement-opsonized HIV-1. In contrast, the bacterial stimulus LPS mediated a non-significant increase in IL-1β at the high concentration in the individual suffering from MWS, while IL-6 was only slightly upregulated by the higher LPS concentration in both the healthy and MWS individual (Figure 1a and 1b).

References

1. Triantafilou K, Hughes TR, Triantafilou M, Morgan BP. The complement membrane attack complex triggers intracellular Ca²⁺ fluxes leading to NLRP3 inflammasome activation. *J Cell Sci.* 2013; 126: 2903-13.
2. Henderson C, Goldbach-Mansky R. Monogenic IL-1 mediated autoinflammatory and immunodeficiency syndromes: finding the right balance in response to danger signals. *Clin Immunol.* 2013; 135: 210-22.
3. Wilflingseder D, Spruth M, Ammann CG, Dopfer S, Speth C, Dierich MP, et al: Complement-mediated enhancement of HIV-1 neutralisation by anti-HLA antibodies derived from polytransfused patients. *Int Arch Allergy Immunol.* 2003; 131: 62-72.
4. Bermejo-Jambrina M, Blatzer M, Jauregui-Onieva P, Yordanov TE, Hortnagl P, Valovka T, et al: CR4 Signaling Contributes to a DC-Driven Enhanced Immune Response Against Complement-Opsonized HIV-1. *Front Immunol.* 2020; 11:1-12.